

# NICOLE WINES LIVESTOCK, LLC



Nicole: 406-220-3329 - Office: 406-224-2099 - nicole@nwlsale.com - nicolewineslivestock.com



## Silver Vacc

Minimum of one round:

- Vaccinate with 7 or 8 Way combination with Somnus
- Vaccinate with IBR, PI3, BVD, BRSV
- Vaccinate with Mannheimia Haemolytica



## Gold Vacc

Two rounds (Spring and Fall):

- Vaccinate w/ 7 or 8 Way at branding, and 7 or 8Way w/ Somnus at preconditioning
- Vaccinate with IBR,PI3,BVD,BRSV
- Vaccinate with one round Mannheimia Haemolytica



## Diamond Vacc

This vaccine tier is for calves weaned 45 days or more

Two rounds ( Spring and Fall):

- Vaccinate w/ 7 or 8 Way at branding, and 7 or 8 Way w/ Somnus at preconditioning
- Vaccinate with IBR,PI3,BVD, and BRSV
- Vaccinate with two rounds Mannheimia Haemolytica
- Internal and/or External parasite control

## Cow/ Bull Vaccine Options

### Yearling Replacement Heifers:

- Pre-breeding: (30-45 days)
  - Vaccinate with a IBR, PI3, BVD, BRSV
  - Vaccinate with Vibrio/Lepto
  - Vaccinate with Trich vaccine (if at risk)
  - Pour-on fly control

### Cows:

- Pre-breeding: (21-30 days)
  - Vaccinate with IBR, PI3, BVD, BRSV
  - Vaccinate with Vibrio/Lepto
  - Vaccinate with Trich (if at risk)

### Bulls:

- Pre-breeding: (30-60 days)
  - Trich test bulls
  - Vaccinate with the same products used on the cows
  - Pour-on for fly control

## Calf Respiratory Vaccine Options

### Branding or out to grass:

- Vaccinate with Blackleg Protection - 7 or 8 way combination vaccines w/ Haemophilus Somnus
- Vaccinate with Pneumonia Protection - IBR,PI3, BVD, BRSV - along with a Mannheimia Haemolytica

### Weaning or Pre-Weaning:

- Booster your Blackleg Protection - 7 or 8 way combination vaccines w/ Haemophilus Somnus
- Booster your Pneumonia Protection
- Booster or vaccinate with a Mannheimia haemolytica

## Pneumonia Protection Options

Ranchers traditionally vaccinate calves around weaning time to reduce "shipping fever" problems. Product administrations, three to six weeks prior to Periods of maximum stress, are encouraged. For the best pneumonia protection, include both viruses (IBR, PI3, BVD and BRSV) and bacteria (Mannheimia Haemolytica) in your program. Our feeders may also recommend, and some even request, vaccines for Haemophilus somnus.

### Modified Live Virus Program

- Option 1 – IBR, PI3, BVD, BRSV
- Option 2 – Viruses + Mannheimia Haemolytica
- Option 3 – Virus + Mannheimia Haemolytica, multocida.  
(Note: add Haemophilus Somnus to Blackleg vaccine.)

### Killed Virus Program

- Option 1 – IBR, PI3, BVD, BRSV
- Option 2 – Viruses + Mannheimia Haemolytica
- Option 3 – Viruses, Mannheimia Haemolytica, Haemophilus somnus

**There are modified live products that can be given to bred cows and calves nursing bred cows when following label directions!**

Modified live virus vaccines and killed virus vaccines each have advantages and disadvantages. Modified vaccines may also include label restrictions. Ask your veterinarian about the best product for your herd, and when it should be administered. In some geographic areas a vitamin E-selenium injection along with vaccine in the fall will enhance disease resistance. Calves to be kept on the home ranch for 40 days or more after weaning will return a profit on de-worming and lice control.